

---

# **Gamma and Poisson Distribution**

## **ECE 341: Random Processes**

### **Lecture #14**

note: All lecture notes, homework sets, and solutions are posted on [www.BisonAcademy.com](http://www.BisonAcademy.com)

---

---

# Gamma and Poisson Distribution

Both are extensions of the exponential distribution:

$$p(t) = ae^{-at}$$

Gamma: Time until  $k$  events occur.

- The time until  $k$  customers arrive,
- The time until  $k$  atoms decay,
- The time until you've been invited to  $k$  parties,

Poisson: Probability  $k$  events occur in  $M$  seconds

- The number of pieces of mail you receive each day (the sending time is exponential)
  - The number of cars through in intersection in one minute
  - The number of customers arriving at a restaurant in one hour
  - Also used to approximate binomial distributions where  $n$  is large
-

---

# Gamma Distribution


Exponential: Time until next event

Gamma: Time until k events

	Exponential	Gamma
pdf	$f_x = a e^{-ax}$	$f_x = \left( \frac{a^k}{(k-1)!} \right) x^{k-1} e^{-ax}$
cdf	$F_x = 1 - e^{-ax}$	complicated
mgf	$\left( \frac{a}{s+a} \right)$	$\left( \frac{a}{s+a} \right)^k$
mean	$\left( \frac{1}{a} \right)$	$\left( \frac{k}{a} \right)$
variance	$\left( \frac{1}{a^2} \right)$	$\left( \frac{k}{a^2} \right)$

---

## pdf of a Gamma distribution:



pdf for a Gamma distribution with an average arrival time of 1

---

## Derivation of pdf from mgf:

Example 1: Determine the pdf and cdf for a Gamma distribution with

- $k = 3$ , and
- $a = 0.2$

Solution: The moment generating function (i.e. LaPlace transform) is

$$\psi(s) = \left( \frac{0.2}{s+0.2} \right)^3$$

From a table of LaPlace transforms (CRC handbook of Mathematics, 1964 edition)

$$\left( \frac{1}{s-a} \right)^3 \rightarrow \frac{1}{2!} t^2 e^{at} u(t)$$

Substituting

$$f_x = (0.2)^3 \frac{1}{2} t^2 e^{-0.2t} u(t)$$

---

---

The cumulative density function (cdf) is the integral of the pdf

$$F_X(s) = \left( \frac{0.2}{s+0.2} \right)^3 \left( \frac{1}{s} \right)$$

Doing partial fraction expansion

$$F_X(s) = \left( \frac{1}{s} \right) + \left( \frac{0.04}{(s+0.2)^3} \right) + \left( \frac{-0.2}{(s+0.2)^2} \right) + \left( \frac{-1}{s+0.2} \right)$$

Taking the inverse LaPlace transform gives you the cdf

$$F_x = (1 + (0.04 t^2 - 0.2 t - 1) e^{-0.2t}) u(t)$$



---

## Poisson Distribution.

The Poisson distribution is actually a discrete probability function.

- Gamma: The time until the  $k$ th customer arrives, ( Gamma )
- Poisson: The probability that  $k$  customers will arrive in a fixed interval

The Poisson distribution is useful if you want to know

- How many cars will go through an intersection in one hour,
  - How many customers will arrive in one hour,
  - How many patients will go to the emergency room in one day, or
  - The number of times your boss will notice you over the course of one week.
  - The probability of a binomial distribution when  $n$  is large
-

---

# Assumptions

(Wikipedia)

- $k$  is the number of times an event occurs in an interval and  $k$  can take values 0, 1, 2, ....
  - The occurrence of one event does not affect the probability that a second event will occur. That is, events occur independently.
  - The average rate at which events occur is independent of any occurrences. For simplicity, this is usually assumed to be constant, but may in practice vary with time.
  - Two events cannot occur at exactly the same instant; instead, at each very small subinterval exactly one event either occurs or does not occur.
-



---

The pdf for a Poisson distribution is (wikipedia)


	Exponential	Poisson
pdf	$f_x = a e^{-ax}$	$f(k, \lambda) = \frac{\lambda^k e^{-\lambda}}{k!}$
cdf	$F_x = 1 - e^{-ax}$	complicated
mgf	$\left(\frac{a}{s+a}\right)$	$\exp(\lambda(z - 1))$
mean	$\left(\frac{1}{a}\right)$	$\lambda$
variance	$\left(\frac{1}{a^2}\right)$	$\lambda$

Here,  $\lambda = 1/a$  for equating exponential and Poisson processes.

---

---

The pdf for a Poisson distribution looks like the following ( $\lambda = 10$  ( $a = 1/10$ ) for illustration purposes).



pdf for a Poisson distribution with  $\lambda = \{4, 8, 12\}$ .  
Note that this is a discrete pdf - so only the integer values of k matter

---

---

## Poisson Approximation for a Binomial Distribution

A Poisson distribution is also a good approximation for a binomial distribution where the number of rolls is large. For this approximation, match the means:

$$\lambda = np$$

**Example 1:** Plot the probability density function for a binomial distribution with

$$n = 100, \quad p = 0.05, \quad \lambda = np = 5$$

Binomial:


$$f_1(x) = \binom{100}{x} (0.05)^x (0.95)^{100-x}$$

Poisson (  $\lambda = np = 5$  )

$$f_2(x) = \frac{1}{x!} \cdot 5^x \cdot e^{-5}$$

---

# Poisson vs. Binomial



Binomial (blue) vs. Poisson (red) with  $np = 5$


---

A Poisson approximation is a slightly more complicated approximation for a binomial distribution than a Normal approximation. It's more accurate however.

- A normal distribution goes from  $-\infty$  to  $+\infty$
- A Poisson distribution is zero for  $k < 0$

In the case of a binomial distribution, you'll never get a negative total.

- A Poisson approximation is slightly more accurate.



---

**Example 2:** Plot the probability density function for a binomial distribution with

$$n = 10,000 \quad p = 0.0005 \quad np = 5$$


This doesn't work well using a Binomial pdf:

$$f(x) = \binom{10,000}{x} (0.0005)^x (0.9995)^{10,000-x}$$

Not a problem with a Poisson approximation

- $np = 5$

$$f(x) \approx \frac{1}{x!} \cdot 5^x \cdot e^{-5}$$



---

## Summary

A Gamma distribution is an exponential distribution

- Where you wait until  $N$  events occur
- The moment generating function is  $\left(\frac{a}{s+a}\right)^N$

A Poisson distribution

- Is an exponential distribution where you count how many events occur over a time interval
- Is a good approximation for a binomial distribution