

---

# Chi-Squared Examples

## ECE 341: Random Processes

### Lecture #26

note: All lecture notes, homework sets, and solutions are posted on [www.BisonAcademy.com](http://www.BisonAcademy.com)

---

---

# Chi-Squared Test

- Is the data consistent with an assumed distribution?

## Procedure

- Collect Data
- Split into N bins
- Compare the expected frequency ( $np$ ) for each bin vs. observed frequency (N)

$$\chi^2 = \sum \left( \frac{(np_i - N_i)^2}{np_i} \right)$$

- Use a chi-squared table to convert the chi-squared score to a probability
-

---

# This Lecture:

- Are world temperatures changing?
  - Does the gain of a transistor have a uniform distribution?
  - Does the gain of a transistor have a normal distribution?
  - Am I psychic?
-

# Are world temperatures changing?


Data:

- The National Oceanic and Atmospheric Administration has been monitoring world temperatures since 1880 (141 years of data)

[https://www.ncdc.noaa.gov/cag/global/time-series/globe/land\\_ocean/p12/12/1880-2020.csv](https://www.ncdc.noaa.gov/cag/global/time-series/globe/land_ocean/p12/12/1880-2020.csv)

Null Hypothesis:

- Temperatures are not changing



# Procedure:

Split the years into 4 bins

- Every 35 years


Split the temperatures into 4 bins

- Hottest 25% to coldest 25%

Null Hypothesis:

- Each 35 year interval's temperatures should be equally distributed among the four percentiles.


## World Temperature Deviation



# Chi-Squared Test

| Years        | Tier | np   | Actual | Chi-Squared    |
|--------------|------|------|--------|----------------|
| 1880<br>-    | 1    | 8.81 | 0      | 8.81           |
|              | 2    | 8.81 | 0      | 8.81           |
| 1915         | 3    | 8.81 | 12     | 1.1551         |
|              | 4    | 8.81 | 23     | 22.8554        |
| 1916<br>-    | 1    | 8.81 | 1      | 6.9235         |
|              | 2    | 8.81 | 9      | 0.0041         |
| 1950         | 3    | 8.81 | 15     | 4.3492         |
|              | 4    | 8.81 | 10     | 0.1607         |
| 1951<br>-    | 1    | 8.81 | 2      | 5.264          |
|              | 2    | 8.81 | 23     | 22.8554        |
| 1985         | 3    | 8.81 | 8      | 0.0745         |
|              | 4    | 8.81 | 2      | 5.264          |
| 1986<br>-    | 1    | 8.81 | 33     | 66.4195        |
|              | 2    | 8.81 | 3      | 3.8316         |
| 2020         | 3    | 8.81 | 0      | 8.81           |
|              | 4    | 8.81 | 0      | 8.81           |
| <b>Total</b> |      |      |        | <b>174.397</b> |

# World Temperature Deviation



# Interpreting the Result

Convert the chi-squared score to a probability

- Chi-squared table
- StatTrek

With 15 degrees of freedom (16 bins), a chi-squared score of 174 corresponds to a probability of at least 0.99995 (rounded to 1)

**I'm at least 99.995% certain that the global temperatures are not uniformly distributed by year (i.e. the temperatures are changing).**

- Enter a value for degrees of freedom.
- Enter a value for one, and only one, of the remaining unshaded text boxes.
- Click the **Calculate** button to compute values for the other text boxes.

Degrees of freedom

Chi-square critical value (CV)

$P(X^2 < 174.397)$


$P(X^2 > 174.397)$

# Is Fargo Getting Warmer?

Data:

- Hector Airport has been measuring the temperature in Fargo since 1942
- High / average / low for each month and year
- <https://www.wunderground.com/history/monthly/us/nd/fargo/KFAR/date/2020-7>
- [http://www.bisonacademy.com/ECE111/Code/Fargo\\_Weather\\_Monthly\\_Avg.txt](http://www.bisonacademy.com/ECE111/Code/Fargo_Weather_Monthly_Avg.txt)

Use the yearly average since 1942






# Procedure

There isn't a lot of data (79 data points).


- Split into 9 bins (should get 8.77 events per bin)
- Split years into 3 intervals
- Split temperature into 3 tiers

Count how many times a given year falls into each bin



# Chi-Squared Test

| Years        | Tier   | np   | Actual | Chi-Squared    |
|--------------|--------|------|--------|----------------|
| 1942 - 1967  | hot    | 8.56 | 5      | 1.4806         |
|              | middle | 8.56 | 10     | 0.2422         |
| 1968 - 1993  | hot    | 8.56 | 11     | 0.6955         |
|              | cold   | 8.56 | 6      | 0.7656         |
| 1994 - 2020  | hot    | 8.56 | 9      | 0.0226         |
|              | cold   | 8.56 | 7      | 0.6955         |
| <b>Total</b> |        |      |        | <b>10.0727</b> |



---

# Interpreting the Result

Convert the chi-squared score to a probability

- Chi-squared table
- StatTrek

With 8 degrees of freedom (9 bins), a chi-squared score of 10.07 corresponds to a probability of at least 0.74

**I'm 74% certain that the temperature in Fargo is changing**

- Enter a value for degrees of freedom.
- Enter a value for one, and only one, of the remaining unshaded text boxes.
- Click the **Calculate** button to compute values for the other text boxes.

|                                |                                    |
|--------------------------------|------------------------------------|
| Degrees of freedom             | <input type="text" value="8"/>     |
| Chi-square critical value (CV) | <input type="text" value="10.07"/> |
| $P(X^2 < 10.07)$               | <input type="text" value="0.74"/>  |
| $P(X^2 > 10.07)$               | <input type="text" value="0.26"/>  |

# Does the gain of a transistor have a uniform distribution?


Each transistor's gain is slightly different.

Does a uniform distribution describe the variability in a transistor's gain?

Is the gain measured consistent with a uniform distribution?

Data:

- Measure the gain of 62 Zetex 1051a transistors
- Sort the gains and plot



# Data Analysis

Null Hypothesis:


- The gain of a Zetex 1051a transistor has a uniform distribution over the range of (600, 1200)

Split this into N regions

- (0, 600)
- (600, 700)
- :
- (1100, 1200),
- (1200, infinity)

Count the number of occurrences in each bin


## Zetex 1051a Transistor



# Chi-Squared Test

| gain         | np    | Actual | Chi-Squared    |
|--------------|-------|--------|----------------|
| >1200        | 0     | 0      | 0              |
| 1100 - 1199  | 10.33 | 1      | 8.4268         |
| 1000 - 1099  | 10.33 | 7      | 1.0735         |
| 900 - 999    | 10.33 | 13     | 0.6901         |
| 800 - 899    | 10.33 | 16     | 3.1122         |
| 700 - 799    | 10.33 | 21     | 11.0212        |
| 600 - 699    | 10.33 | 4      | 3.8789         |
| 0 - 599      | 0     | 0      | 0              |
| <b>Total</b> |       |        | <b>28.2027</b> |

## Zetex 1051a Transistor



# Interpreting the Results

Convert the chi-squared score to a probability

- Chi-squared table
- StatTrek

With 7 degrees of freedom (8 bins), a chi-squared score of 28.2 corresponds to a probability of at least 0.9998

**I'm 99.98% certain that the gain of a Zetex 1051a transistor does not have a uniform distribution**

- The data is inconsistent with a uniform distribution

- Enter a value for degrees of freedom.
- Enter a value for one, and only one, of the remaining unshaded text boxes.
- Click the **Calculate** button to compute values for the other text boxes.

|                                |                                     |
|--------------------------------|-------------------------------------|
| Degrees of freedom             | <input type="text" value="7"/>      |
| Chi-square critical value (CV) | <input type="text" value="28.2"/>   |
| $P(X^2 < 28.2)$                | <input type="text" value="0.9998"/> |
| $P(X^2 > 28.2)$                | <input type="text" value="0.0002"/> |


# Does the gain of a transistor have a Normal distribution?

- mean = 854.1290
- standard deviation = 120.2034

Same procedure as before but the probabilities change

- Use a normal distribution and a z-score to determine the probability of each region

**Zetex 1051a Transistor**





# Probabilities of Each Region

- Use StatTrek to find the cdf
- From that, find the probability of each region

| region | cdf   | p(region) |
|--------|-------|-----------|
| 1,200  | 0.998 | 0.018     |
| 1,100  | 0.98  | 0.092     |
| 1,000  | 0.888 | 0.239     |
| 900    | 0.649 | 0.323     |
| 800    | 0.326 | 0.226     |
| 700    | 0.1   | 0.083     |
| 600    | 0.017 | 0.017     |

- Enter a value in three of the four text boxes.
- Leave the fourth text box blank.
- Click the **Calculate** button to compute a value for the blank text box.

Normal random variable (x)

Cumulative probability:  $P(X \leq 800)$

Mean

Standard deviation


# Chi-Squared Calculations

Use the probabilities from the previous slide

| gain         | p     | np     | Actual | Chi-Squared   |
|--------------|-------|--------|--------|---------------|
| >1200        | 0.002 | 0.124  | 0      | 0             |
| 1100 - 1199  | 0.018 | 1.116  | 1      | 0.0121        |
| 1000 - 1099  | 0.092 | 5.704  | 7      | 0.2945        |
| 900 - 999    | 0.239 | 14.818 | 13     | 0.223         |
| 800 - 899    | 0.323 | 20.026 | 16     | 0.8094        |
| 700 - 799    | 0.226 | 14.012 | 21     | 3.485         |
| 600 - 699    | 0.083 | 5.146  | 4      | 0.2552        |
| 0 - 599      | 0.017 | 1.054  | 0      | 1.054         |
| <b>Total</b> |       |        |        | <b>6.1332</b> |

## Zetex 1051a Transistor

Gain (hfe)



# Interpreting the Results

A chi-squared score of 6.13 corresponds to a probability of 0.48

- There is a 48% chance of rejecting the null hypothesis (this is a normal distribution)

Midrange numbers like this mean "no conclusion"

- The data is consistent with a normal distribution
  - *the chi-squared score is not too large*
- It does not appear that the data was fudged
  - *The chi-squared score is not too small*


- Enter a value for degrees of freedom.
- Enter a value for one, and only one, of the remaining unshaded text boxes.
- Click the **Calculate** button to compute values for the other text boxes.

|                                |                                   |
|--------------------------------|-----------------------------------|
| Degrees of freedom             | <input type="text" value="7"/>    |
| Chi-square critical value (CV) | <input type="text" value="6.13"/> |
| $P(X^2 < 6.13)$                | <input type="text" value="0.48"/> |
| $P(X^2 > 6.13)$                | <input type="text" value="0.52"/> |

---

# Am I Psychic?

- Take a deck of playing cards
- Shuffle them
- Predict the suit for the top card
- Flip it up and place in one pile if correct, another pile if incorrect
- Count how many times I'm right
- Use a chi-squared test to see if I'm able to foresee the suit with odds that pure chance cannot explain



# Data

- Predicted Correctly: 10 times
- Predicted Incorrectly: 42 times

## Chi-Squared Test

| case         | np | Actual | Chi-Squared   |
|--------------|----|--------|---------------|
| Correct      | 13 | 10     | 0.6923        |
| Incorrect    | 39 | 42     | 0.2308        |
| <b>Total</b> |    |        | <b>0.9231</b> |

## Result:

- probability = 66%
- There is a 66% chance of rejecting the null hypothesis
  - 66% chance I'm not just guessing randomly
  - 66% chance I'm worse than the monkey score

- Enter a value for degrees of freedom.
- Enter a value for one, and only one, of the remaining unshaded text boxes.
- Click the **Calculate** button to compute values for the other text boxes.

Degrees of freedom

Chi-square critical value (CV)

$P(X^2 < 0.9231)$

$P(X^2 > 0.9231)$